

Chromosome 15

Description

Humans normally have 46 chromosomes in each cell, divided into 23 pairs. Two copies of chromosome 15, one copy inherited from each parent, form one of the pairs. Chromosome 15 spans approximately 102 million DNA building blocks (nucleotides) and represents more than 3 percent of the total DNA in cells.

Researchers are continually improving their methods of genome sequencing and working to confirm the number of genes on each chromosome. Chromosome 15 is estimated to contain around 630 genes that provide instructions for making proteins. These proteins perform a variety of different roles in the body.

Health Conditions Related to Chromosomal Changes

The following chromosomal conditions are associated with changes in the structure or number of copies of chromosome 15.

15q11-q13 duplication syndrome

Duplication of a region of the long (q) arm of chromosome 15 can result in 15q11-q13 duplication syndrome (also called dup15q syndrome). The features of this condition can include weak muscle tone (hypotonia); intellectual disabilities; recurrent seizures (epilepsy); and characteristics typical of autism spectrum disorder, a developmental condition that affects communication and social interaction.

Dup15q syndrome is caused by the presence of at least one extra copy of a region of chromosome 15 called 15q11.2-q13.1. This region is also called the Prader-Willi/Angelman critical region (PWACR) because genetic changes in it are also involved in conditions called Prader-Willi syndrome and Angelman syndrome (described below). Individuals with a 15q11.2-q13.1 duplication will develop the signs and symptoms of dup15q syndrome only if the chromosome abnormality occurs on the copy of the chromosome that comes from the egg cell. During fertilization, the egg and the sperm each contribute one copy of each chromosome. However, some genes on chromosome 15, including some of those in the 15q11.2-q13.1 region, are turned on (active) only on

the copy that came from the egg cell. This parent-specific gene activation results from a phenomenon called genomic imprinting.

The most common chromosome abnormality that leads to 15q11.2-q13.1 duplication, occurring in about 80 percent of people with dup15q syndrome, is called an isodicentric chromosome 15. An isodicentric chromosome contains mirror-image segments of genetic material and has two constriction points (centromeres), rather than the single centromere found in normal chromosomes. In people with an isodicentric chromosome 15, cells have the usual two copies of chromosome 15 plus the two duplicated copies of the segment of genetic material in the isodicentric chromosome, for a total of four copies of the duplicated segment.

In about 20 percent of cases of dup15q syndrome, the duplication occurs on the q arm of one of the two copies of chromosome 15 in each cell; this situation is called an interstitial duplication. In these cases, cells have two copies of chromosome 15, one of which has an extra copy of the segment of genetic material, for a total of three copies of the duplicated segment.

In all cases of dup15q syndrome, the duplicated genetic material results in extra copies of certain genes involved in development. This extra genetic material disrupts normal development, causing the characteristic features of this disorder. People with dup15q syndrome caused by an interstitial duplication often have milder signs and symptoms than those with an isodicentric chromosome 15.

15q13.3 microdeletion

A 15q13.3 microdeletion is a chromosomal change in which a small piece of chromosome 15 is deleted in each cell. The deletion occurs on the q arm of the chromosome at a position designated q13.3. Most people with a 15q13.3 microdeletion are missing a sequence of about 2 million DNA nucleotides, also written as 2 megabases (Mb). The exact size of the deleted region varies, but it typically contains at least six genes. It is unclear how a loss of these genes increases the risk of intellectual disabilities, seizures, behavioral problems, and psychiatric disorders in some individuals with a 15q13.3 microdeletion.

Other people with a 15q13.3 microdeletion have no obvious signs or symptoms related to the chromosomal change. In these individuals, the microdeletion is often detected when they undergo genetic testing because they have an affected relative. It is unknown why a 15q13.3 microdeletion causes cognitive and behavioral problems in some individuals but few or no health problems in others. Researchers believe that additional genetic or environmental factors may be involved.

15q24 microdeletion

A 15q24 microdeletion is a chromosomal change in which a small piece of chromosome 15 is deleted in each cell. Specifically, affected individuals are missing between 1.7 Mb and 6.1 Mb of DNA at position q24 on chromosome 15. The exact size of the deletion varies, but all individuals are missing the same 1.2 Mb region. This region contains several genes that are thought to be important for normal development. It is unclear

how a loss of these genes leads to the intellectual disabilities, distinctive facial features, and other abnormalities often seen in people with a 15q24 microdeletion.

Acute promyelocytic leukemia

A type of blood cancer known as acute promyelocytic leukemia is caused by a rearrangement (translocation) of genetic material between chromosomes 15 and 17. This translocation, written as t(15;17), fuses part of the *PML* gene from chromosome 15 with part of the *RARA* gene from chromosome 17. This variant (also called a mutation) is acquired during a person's lifetime and is present only in certain cells. This type of genetic change, called a somatic variant, is not inherited. The t(15;17) translocation is called a balanced reciprocal translocation because the pieces of chromosome are exchanged with each other (reciprocal) and no genetic material is gained or lost (balanced). The protein produced from this fused gene is known as PML-RAR α .

The PML-RAR α protein functions differently than the proteins produced from the normal *PML* and *RARA* genes. The *PML* gene on chromosome 15 provides instructions for a protein that acts as a tumor suppressor, which means it prevents cells from growing and dividing too rapidly or in an uncontrolled way. The PML protein blocks cell growth and division (proliferation) and induces self-destruction (apoptosis) in combination with other proteins.

The *RARA* gene on chromosome 17 provides instructions for making a transcription factor called the retinoic acid receptor alpha (RAR α). A transcription factor is a protein that attaches (binds) to specific regions of DNA and helps control the activity of particular genes. Normally, the RAR α protein controls the activity of genes that are important for the maturation (differentiation) of immature white blood cells beyond a particular stage called the promyelocyte.

The PML-RAR α protein interferes with the normal function of both the PML and the RAR α proteins. As a result, blood cells are stuck at the promyelocyte stage, and they proliferate abnormally. Excess promyelocytes accumulate in the bone marrow, and normal white blood cells cannot form, leading to acute promyelocytic leukemia.

Angelman syndrome

Angelman syndrome is a condition that results from a loss of gene activity (expression) in a specific part of chromosome 15 in each cell. This region is located on the q arm of the chromosome and is designated 15q11-q13. This region contains a gene called *UBE3A*. The neurological features of Angelman syndrome are caused by genetic changes that affect this gene.

Normally, both the egg and sperm cells each contribute one copy of the *UBE3A* gene during fertilization, and both copies of this gene are active in most of the body's tissues. However, in nerve cells (neurons) in the brain and spinal cord, only the copy derived from the egg cell is active. If this copy is lost because of a chromosomal change or a gene variant, a person will have no working copies of the *UBE3A* gene in some parts of the brain.

In most cases (about 70 percent), Angelman syndrome is a result of a deletion in the copy of chromosome 15 derived from the egg cell. This chromosomal change deletes the region of chromosome 15 that includes the *UBE3A* gene. Because the copy of the *UBE3A* gene derived from the sperm cell is normally inactive in certain parts of the brain, a deletion in the copy of chromosome 15 from the egg cell leaves no active copies of the *UBE3A* gene in these brain regions.

In 3 percent to 7 percent of cases of Angelman syndrome, the condition occurs when a person has two copies of chromosome 15 derived from the sperm cell instead of one copy from the egg and sperm cell. This phenomenon is called uniparental disomy (UPD). People with UPD for chromosome 15 have two copies of the *UBE3A* gene, but because both of those copies are derived from sperm cells, they are therefore inactive in the brain.

About 10 percent of cases of Angelman syndrome are caused by a variant in the *UBE3A* gene, and another 3 percent are caused by a defect in the DNA region that controls the activation of the *UBE3A* gene and other genes on the copy of chromosome 15 that is derived from egg cells. In a small percentage of cases, Angelman syndrome is caused by a translocation or by a variant in a gene other than *UBE3A*. These genetic changes abnormally inactivate the *UBE3A* gene.

Deafness-infertility syndrome

Deafness-infertility syndrome is caused by a deletion of genetic material on the q arm of chromosome 15. The symptoms of deafness-infertility syndrome are related to the loss of multiple genes in this region. The size of the deletion varies among affected individuals. Researchers have determined that the loss of a particular gene on chromosome 15, *STRC*, is responsible for hearing loss in affected individuals. The loss of another gene, *CATSPER2*, in the same region of chromosome 15, can lead to abnormalities in sperm. People with this deletion are unable to conceive children (a condition called infertility). Researchers are working to determine how the loss of additional genes in the deleted region affects people with deafness-infertility syndrome.

Prader-Willi syndrome

Prader-Willi syndrome is caused by a loss of active genes in a region of chromosome 15. This region is located on the q arm of the chromosome and is designated 15q11-q13. It is the same part of chromosome 15 that is usually affected in people with Angelman syndrome, although different genes are associated with the two disorders. People can have either Prader-Willi syndrome or Angelman syndrome, but they typically cannot have both.

Normally, both the egg and sperm cells each contribute one copy of chromosome 15 during fertilization. Some genes on this chromosome are active only on the copy from the sperm cell. In about 70 percent of cases, Prader-Willi syndrome occurs when the 15q11-q13 region in the copy of chromosome 15 that is derived from the sperm cell is deleted. A person with this chromosomal change will be missing certain critical genes in this region because the genes on the copy from the egg cell are turned off (inactive).

Researchers are working to identify which missing genes are associated with the characteristic features of Prader-Willi syndrome.

In about 25 percent of cases, people with Prader-Willi syndrome have two copies of chromosome 15 derived from the egg cell instead of one copy from the egg and sperm cell. This phenomenon is called UPD. A person with two copies of chromosome 15 derived from the egg cell will have no active copies of certain genes in the 15q11-q13 region.

In a small percentage of cases, Prader-Willi syndrome is caused by a translocation. Rarely, the condition results from a variant or other change that abnormally inactivates genes on the copy of chromosome 15 derived from the sperm cell.

Other chromosomal conditions

Other changes in the structure or number of copies of chromosome 15 can cause intellectual disabilities, delayed growth and development, hypotonia, and characteristic facial features. These changes include an extra copy of part of chromosome 15 in each cell (partial trisomy 15), a missing segment of the chromosome in each cell (partial monosomy 15), and a circular structure called ring chromosome 15. A ring chromosome occurs when a chromosome breaks in two places and the ends of the chromosome arms fuse together to form a circular structure.

Additional Information & Resources

Additional NIH Resources

- National Human Genome Research Institute: Chromosome Abnormalities (<https://www.genome.gov/about-genomics/fact-sheets/Chromosome-Abnormalities-Fact-Sheet>)

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28Chromosomes,+Human,+Pair+15%5BMAJR%5D%29+AND+%28Chromosome+15%5BTI%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1440+days%22%5Bdp%5D>)

References

- Battaglia A. The inv dup (15) or idic (15) syndrome (Tetrasomy 15q). *Orphanet J Rare Dis.* 2008 Nov 19;3:30. doi: 10.1186/1750-1172-3-30. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19019226>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2613132/>)
- Bittel DC, Butler MG. Prader-Willi syndrome: clinical genetics, cytogenetics and molecular biology. *Expert Rev Mol Med.* 2005 Jul 25;7(14):1-20. doi:10.1017/S1462399405009531. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/160386>)

20)

- Bittel DC, Kibiryeva N, Talebizadeh Z, Butler MG. Microarray analysis of gene/transcript expression in Prader-Willi syndrome: deletion versus UPD. *J Med Genet*. 2003 Aug;40(8):568-74. doi: 10.1136/jmg.40.8.568. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12920063>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1735542/>)
- Bittel DC, Kibiryeva N, Talebizadeh Z, Driscoll DJ, Butler MG. Microarray analysis of gene/transcript expression in Angelman syndrome: deletion versus UPD. *Genomics*. 2005 Jan;85(1):85-91. doi: 10.1016/j.ygeno.2004.10.010. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15607424>)
- Borgatti R, Piccinelli P, Passoni D, Dalpra L, Miozzo M, Micheli R, Gagliardi C, Balottin U. Relationship between clinical and genetic features in inverted duplicated chromosome 15 patients. *Pediatr Neurol*. 2001 Feb;24(2):111-6. doi:10.1016/s0887-8994(00)00244-7. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11275459>)
- Buiting K. Prader-Willi syndrome and Angelman syndrome. *Am J Med Genet C Semin Med Genet*. 2010 Aug 15;154C(3):365-76. doi: 10.1002/ajmg.c.30273. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20803659>)
- Butler MG, Bittel DC, Kibiryeva N, Talebizadeh Z, Thompson T. Behavioral differences among subjects with Prader-Willi syndrome and type I or type II deletion and maternal disomy. *Pediatrics*. 2004 Mar;113(3 Pt 1):565-73. doi:10.1542/peds.113.3.565. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/14993551>)
- Cassidy SB, Dykens E, Williams CA. Prader-Willi and Angelman syndromes: sister imprinted disorders. *Am J Med Genet*. 2000 Summer;97(2):136-46. doi:10.1002/1096-8628(200022)97:2<136::AID-AMJG13>3.0.CO;2-V. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11180221>)
- Clayton-Smith J, Laan L. Angelman syndrome: a review of the clinical and genetic aspects. *J Med Genet*. 2003 Feb;40(2):87-95. doi: 10.1136/jmg.40.2.87. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12566516>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1735357/>)
- Collins SJ. The role of retinoids and retinoic acid receptors in normal hematopoiesis. *Leukemia*. 2002 Oct;16(10):1896-905. doi: 10.1038/sj.leu.2402718. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12357341>)
- de The H, Lavau C, Marchio A, Chomienne C, Degos L, Dejean A. The PML-RAR α fusion mRNA generated by the t(15;17) translocation in acute promyelocytic leukemia encodes a functionally altered RAR. *Cell*. 1991 Aug 23;66(4):675-84. doi:10.1016/0092-8674(91)90113-d. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/1652369>)
- El-Hattab AW, Zhang F, Maxim R, Christensen KM, Ward JC, Hines-Dowell S, Scaglia F, Lupski JR, Cheung SW. Deletion and duplication of 15q24: molecular mechanisms and potential modification by additional copy number variants. *Genet Med*. 2010 Sep;12(9):573-86. doi: 10.1097/GIM.0b013e3181eb9b4a. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20860070>)

- Ensembl Human Map View (http://www.ensembl.org/Homo_sapiens/Location/Chromosome?chr=15;r=15:1-101991189)
- Gilbert F. Disease genes and chromosomes: disease maps of the human genome. *Chromosome 15. Genet Test.* 1999;3(3):309-22. doi: 10.1089/109065799316653. Noabstract available. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10495933>)
- Horsthemke B, Buiting K. Imprinting defects on human chromosome 15. *CytogenetGenome Res.* 2006;113(1-4):292-9. doi: 10.1159/000090844. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16575192>)
- Horsthemke B, Wagstaff J. Mechanisms of imprinting of thePrader-Willi/Angelman region. *Am J Med Genet A.* 2008 Aug 15;146A(16):2041-52.doi: 10.1002/ajmg.a.32364. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18627066>)
- Lee S, Wevrick R. Identification of novel imprinted transcripts in thePrader-Willi syndrome and Angelman syndrome deletion region: further evidence forregional imprinting control. *Am J Hum Genet.* 2000 Mar;66(3):848-58. doi:10.1086/302817. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10712201>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1288168/>)
- Pandolfi PP. Oncogenes and tumor suppressors in the molecular pathogenesis ofacute promyelocytic leukemia. *Hum Mol Genet.* 2001 Apr;10(7):769-75. doi:10.1093/hmg/10.7.769. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11257111>)
- Rineer S, Finucane B, Simon EW. Autistic symptoms among children and youngadults with isodicentric chromosome 15. *Am J Med Genet.* 1998 Sep 7;81(5):428-33.doi: 10.1002/(sici)1096-8628(19980907)81:53.0.co;2-e. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9754629>)
- Salomoni P, Pandolfi PP. The role of PML in tumor suppression. *Cell.* 2002 Jan25;108(2):165-70. doi: 10.1016/s0092-8674(02)00626-8. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11832207>)
- Sharp AJ, Mefford HC, Li K, Baker C, Skinner C, Stevenson RE, Schroer RJ, Novara F, De Gregori M, Ciccone R, Broomer A, Casuga I, Wang Y, Xiao C, Barbacioru C, Gimelli G, Bernardina BD, Torniero C, Giorda R, Regan R, Murday V, Mansour S, Fichera M, Castiglia L, Failla P, Ventura M, Jiang Z, Cooper GM, Knight SJ, Romano C, Zuffardi O, Chen C, Schwartz CE, Eichler EE. A recurrent15q13.3 microdeletion syndrome associated with mental retardation and seizures.*Nat Genet.* 2008 Mar;40(3):322-8. doi: 10.1038/ng.93. Epub 2008 Feb 17. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18278044>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2365467/>)
- Sharp AJ, Selzer RR, Veltman JA, Gimelli S, Gimelli G, Striano P, Coppola A, Regan R, Price SM, Knoers NV, Eis PS, Brunner HG, Hennekam RC, Knight SJ, deVries BB, Zuffardi O, Eichler EE. Characterization of a recurrent 15q24microdeletion syndrome. *Hum Mol Genet.* 2007 Mar 1;16(5):567-72. doi:10.1093/hmg/ddm016. Epub 2007 Mar 14. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17360722>)
- Tumer Z, Harboe TL, Blennow E, Kalscheuer VM, Tommerup N, Brondum-Nielsen

K.Molecular cytogenetic characterization of ring chromosome 15 in three unrelated patients. *Am J Med Genet A*. 2004 Nov 1;130A(4):340-4. doi: 10.1002/ajmg.a.30035. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15384084>)

- UCSC Genome Browser: Statistics (<http://genome.cse.ucsc.edu/goldenPath/stats.html>)
- van Bon BW, Mefford HC, Menten B, Koolen DA, Sharp AJ, Nillesen WM, Innis JW, de Ravel TJ, Mercer CL, Fichera M, Stewart H, Connell LE, Ounap K, Lachlan K, Castle B, Van der Aa N, van Ravenswaaij C, Nobrega MA, Serra-Juhe C, Simonic I, de Leeuw N, Pfundt R, Bongers EM, Baker C, Finnemore P, Huang S, Maloney VK, Crolla JA, van Kalmthout M, Elia M, Vandeweyer G, Fryns JP, Janssens S, Foulds N, Reitano S, Smith K, Parkel S, Loeys B, Woods CG, Oostra A, Speleman F, Pereira AC, Kurg A, Willatt L, Knight SJ, Vermeesch JR, Romano C, Barber JC, Mortier G, Perez-Jurado LA, Kooy F, Brunner HG, Eichler EE, Kleefstra T, de Vries BB. Further delineation of the 15q13 microdeletion and duplication syndromes: a clinical spectrum varying from non-pathogenic to a severe outcome. *J Med Genet*. 2009 Aug;46(8):511-23. doi: 10.1136/jmg.2008.063412. Epub 2009 Apr 15. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19372089>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3395372/>)
- Zhang Y, Malekpour M, Al-Madani N, Kahrizi K, Zanganeh M, Lohr NJ, Mohseni M, Mojahedi F, Daneshi A, Najmabadi H, Smith RJ. Sensorineural deafness and male infertility: a contiguous gene deletion syndrome. *J Med Genet*. 2007 Apr;44(4):233-40. doi: 10.1136/jmg.2006.045765. Epub 2006 Nov 10. Erratum In: *J Med Genet*. 2007 Aug;44(8):544. Lohr, Naomi J [added]. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17098888>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2598039/>)
- Zody MC, Garber M, Sharpe T, Young SK, Rowen L, O'Neill K, Whittaker CA, Kamal M, Chang JL, Cuomo CA, Dewar K, FitzGerald MG, Kodira CD, Madan A, Qin S, Yang X, Abbasi N, Abouelleil A, Arachchi HM, Baradarani L, Birditt B, Bloom S, Bloom T, Borowsky ML, Burke J, Butler J, Cook A, DeArellano K, DeCaprio D, Dorris L 3rd, Dors M, Eichler EE, Engels R, Fahey J, Fleetwood P, Friedman C, Gearin G, Hall JL, Hensley G, Johnson E, Jones C, Kamat A, Kaur A, Locke DP, Madan A, Munson G, Jaffe DB, Lui A, Macdonald P, Mauceli E, Naylor JW, Nesbitt R, Nicol R, O'Leary SB, Ratcliffe A, Rounsley S, She X, Sneddon KM, Stewart S, Sougnez C, Stone SM, Topham K, Vincent D, Wang S, Zimmer AR, Birren BW, Hood L, Lander ES, Nusbaum C. Analysis of the DNA sequence and duplication history of human chromosome 15. *Nature*. 2006 Mar 30;440(7084):671-5. doi: 10.1038/nature04601. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16572171>)
- Zollino M, Tiziano F, Di Stefano C, Neri G. Partial duplication of the long arm of chromosome 15: confirmation of a causative role in craniosynostosis and definition of a 15q25-qter trisomy syndrome. *Am J Med Genet*. 1999 Dec 22;87(5):391-4. doi: 10.1002/(sici)1096-8628(19991222)87:53.0.co;2-o. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10594876>)

Last updated April 26, 2024