

## **CLN7** disease

## **Description**

CLN7 disease is an inherited disorder that primarily affects the nervous system. The signs and symptoms of this condition typically begin between ages 2 and 7. The initial features are usually vision loss and problems with movement that might seem like clumsiness. Additional signs and symptoms of CLN7 disease include muscle twitches (myoclonus), difficulty coordinating movements (ataxia), recurrent seizures (epilepsy), and speech impairment. Mental functioning and motor skills (such as sitting and walking) decline with age. Individuals with CLN7 disease typically do not survive past their teens.

CLN7 disease is one of a group of disorders known as neuronal ceroid lipofuscinoses ( NCLs), which may also be collectively referred to as Batten disease. All these disorders affect the nervous system and typically cause worsening problems with vision, movement, and thinking ability. The different NCLs are distinguished by their genetic cause. Each disease type is given the designation "CLN," meaning ceroid lipofuscinosis, neuronal, and then a number to indicate its subtype.

# Frequency

The incidence of CLN7 disease is unknown; more than 70 cases have been described in the scientific literature. CLN7 disease was first diagnosed in the Turkish population and was thought to be limited to individuals in that group. However, CLN7 disease has now been identified in people around the world. Collectively, all forms of NCL affect an estimated 1 in 100,000 individuals worldwide.

#### Causes

Mutations in the *MFSD8* gene cause CLN7 disease. The *MFSD8* gene provides instructions for making a protein whose function is unknown. The MFSD8 protein is embedded in the membrane of cell compartments called lysosomes, which digest and recycle different types of molecules. Based on the structure of the protein, MFSD8 probably transports molecules across the lysosomal membrane, but the specific molecules it moves have not been identified.

*MFSD8* gene mutations likely lead to the production of a protein with altered structure or function. It is unclear how an altered MFSD8 protein leads to the severe neurological

features of CLN7 disease. CLN7 disease, like other NCLs, is characterized by the accumulation of proteins and other substances in lysosomes. These accumulations occur in cells throughout the body; however, nerve cells seem to be particularly vulnerable to their effects. These accumulations can cause cell damage leading to cell death. Individuals with CLN7 disease have gradual nerve cell loss in certain parts of the brain, which likely leads to the signs and symptoms of this condition.

# Learn more about the gene associated with CLN7 disease

MFSD8

### Inheritance

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

#### Other Names for This Condition

- CLN7
- CLN7 disease, late infantile
- MFSD8-related neuronal ceroid lipofuscinosis

### **Additional Information & Resources**

#### Genetic Testing Information

Genetic Testing Registry: Neuronal ceroid lipofuscinosis 7 (https://www.ncbi.nlm.nih.gov/gtr/conditions/C1838571/)

## Genetic and Rare Diseases Information Center

CLN7 disease (https://rarediseases.info.nih.gov/diseases/1220/index)

#### Patient Support and Advocacy Resources

National Organization for Rare Disorders (NORD) (https://rarediseases.org/)

#### Clinical Trials

ClinicalTrials.gov (https://clinicaltrials.gov/search?cond=%22CLN7 disease%22)

## Catalog of Genes and Diseases from OMIM

• CEROID LIPOFUSCINOSIS, NEURONAL, 7; CLN7 (https://omim.org/entry/610951)

## Scientific Articles on PubMed

PubMed (https://pubmed.ncbi.nlm.nih.gov/?term=%28%28CLN7+disease%5BTIAB %5D%29+OR+%28CLN7%5BTIAB%5D%29+AND+%28neuronal+ceroid+lipofuscin osis%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+A ND+%22last+3600+days%22%5Bdp%5D)

#### References

- Craiu D, Dragostin O, Dica A, Hoffman-Zacharska D, Gos M, Bastian AE, Gherghiceanu M, Rolfs A, Nahavandi N, Craiu M, Iliescu C. Rett-like onset inlate-infantile neuronal ceroid lipofuscinosis (CLN7) caused by compoundheterozygous mutation in the MFSD8 gene and review of the literature data onclinical onset signs. Eur J Paediatr Neurol. 2015 Jan;19(1):78-86. doi:10.1016/j.ejpn.2014.07.008. Epub 2014 Aug 7. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/25439737)
- Kollmann K, Uusi-Rauva K, Scifo E, Tyynela J, Jalanko A, Braulke T. Cellbiology and function of neuronal ceroid lipofuscinosis-related proteins. BiochimBiophys Acta. 2013 Nov;1832(11):1866-81. doi: 10.1016/j.bbadis.2013.01.019. Epub2013 Feb 9. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/23402926)
- Kousi M, Siintola E, Dvorakova L, Vlaskova H, Turnbull J, Topcu M, Yuksel D, Gokben S, Minassian BA, Elleder M, Mole SE, Lehesjoki AE. Mutations in CLN7/ MFSD8are a common cause of variant late-infantile neuronal ceroid lipofuscinosis. Brain. 2009 Mar;132(Pt 3):810-9. doi: 10.1093/brain/awn366. Epub 2009 Feb 5. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/19201763)
- Sharifi A, Kousi M, Sagne C, Bellenchi GC, Morel L, Darmon M, Hulkova H, RuivoR, Debacker C, El Mestikawy S, Elleder M, Lehesjoki AE, Jalanko A, Gasnier B,Kyttala A. Expression and lysosomal targeting of CLN7, a major facilitatorsuperfamily transporter associated with variant late-infantile neuronal ceroidlipofuscinosis. Hum Mol Genet. 2010 Nov 15;19(22):4497-514. doi:10.1093/hmg/ddq381. Epub 2010 Sep 7. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/20826447) or Free article on PubMed Central (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3298853/)
- Siintola E, Topcu M, Aula N, Lohi H, Minassian BA, Paterson AD, Liu XQ, WilsonC, Lahtinen U, Anttonen AK, Lehesjoki AE. The novel neuronal ceroidlipofuscinosis gene MFSD8 encodes a putative lysosomal transporter. Am J HumGenet. 2007 Jul; 81(1):136-46. doi: 10.1086/518902. Epub 2007 May 14. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/17564970) or Free article on PubMed Central (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1950917/)
- Williams RE, Mole SE. New nomenclature and classification scheme for theneuronal ceroid lipofuscinoses. Neurology. 2012 Jul 10;79(2):183-91. doi:10.1212/

WNL.0b013e31825f0547. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/22778232)

Last updated June 1, 2021