

Fish-eye disease

Description

Fish-eye disease, also called partial LCAT deficiency, is a disorder that causes the clear front surface of the eyes (the corneas) to gradually become cloudy. The cloudiness, which generally first appears in adolescence or early adulthood, consists of small grayish dots of cholesterol (opacities) distributed across the corneas. Cholesterol is a waxy, fat-like substance that is produced in the body and obtained from foods that come from animals; it aids in many functions of the body but can become harmful in excessive amounts. As fish-eye disease progresses, the corneal cloudiness worsens and can lead to severely impaired vision.

Frequency

Fish-eye disease is a rare disorder. Approximately 30 cases have been reported in the medical literature.

Causes

Fish-eye disease is caused by mutations in the *LCAT* gene. This gene provides instructions for making an enzyme called lecithin-cholesterol acyltransferase (LCAT).

The LCAT enzyme plays a role in removing cholesterol from the blood and tissues by helping it attach to molecules called lipoproteins, which carry it to the liver. Once in the liver, the cholesterol is redistributed to other tissues or removed from the body. The enzyme has two major functions, called alpha- and beta-LCAT activity. Alpha-LCAT activity helps attach cholesterol to a lipoprotein called high-density lipoprotein (HDL). Beta-LCAT activity helps attach cholesterol to other lipoproteins called very low-density lipoprotein (VLDL) and low-density lipoprotein (LDL).

LCAT gene mutations that cause fish-eye disease impair alpha-LCAT activity, reducing the enzyme's ability to attach cholesterol to HDL. Impairment of this mechanism for reducing cholesterol in the body leads to cholesterol-containing opacities in the corneas. It is not known why the cholesterol deposits affect only the corneas in this disorder. Mutations that affect both alpha-LCAT activity and beta-LCAT activity lead to a related disorder called complete LCAT deficiency, which involves corneal opacities in combination with features affecting other parts of the body.

Learn more about the gene associated with Fish-eye disease

LCAT

Inheritance

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- Alpha-LCAT deficiency
- Alpha-lecithin:cholesterol acyltransferase deficiency
- Dyslipoproteinemic corneal dystrophy
- FED
- LCATA deficiency
- Partial LCAT deficiency

Additional Information & Resources

Genetic Testing Information

Genetic Testing Registry: Fish-eye disease (https://www.ncbi.nlm.nih.gov/gtr/conditions/C0342895/)

Genetic and Rare Diseases Information Center

Fish-eye disease (https://rarediseases.info.nih.gov/diseases/6450/index)

Patient Support and Advocacy Resources

National Organization for Rare Disorders (NORD) (https://rarediseases.org/)

Clinical Trials

ClinicalTrials.gov (https://clinicaltrials.gov/search?cond=%22Fish-eye disease%22)

Catalog of Genes and Diseases from OMIM

FISH-EYE DISEASE; FED (https://omim.org/entry/136120)

Scientific Articles on PubMed

 PubMed (https://pubmed.ncbi.nlm.nih.gov/?term=%28%28fish-eye+disease%5BTIA B%5D%29+OR+%28alpha-lcat+deficiency%5BTIAB%5D%29+OR+%28cholesterol+ acyltransferase+deficiency%5BTIAB%5D%29+OR+%28partial+lcat+deficiency%5B TIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22l ast+3600+days%22%5Bdp%5D)

References

- Calabresi L, Pisciotta L, Costantin A, Frigerio I, Eberini I, Alessandrini P,Arca M, Bon GB, Boscutti G, Busnach G, Frasca G, Gesualdo L, Gigante M,Lupattelli G, Montali A, Pizzolitto S, Rabbone I, Rolleri M, Ruotolo G, Sampietro T, Sessa A, Vaudo G, Cantafora A, Veglia F, Calandra S, Bertolini S, Franceschini G. The molecular basis of lecithin:cholesterol acyltransferase deficiencysyndromes: a comprehensive study of molecular and biochemical findings in 13unrelated Italian families. Arterioscler Thromb Vasc Biol. 2005 Sep;25(9):1972-8.doi: 10.1161/01.ATV. 0000175751.30616.13. Epub 2005 Jun 30. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/15994445)
- Contacos C, Sullivan DR, Rye KA, Funke H, Assmann G. A new molecular defect inthe lecithin: cholesterol acyltransferase (LCAT) gene associated with fish eyedisease. J Lipid Res. 1996 Jan;37(1):35-44. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/8820100)
- Kuivenhoven JA, Stalenhoef AF, Hill JS, Demacker PN, Errami A, Kastelein JJ, Pritchard PH. Two novel molecular defects in the LCAT gene are associated withfish eye disease. Arterioscler Thromb Vasc Biol. 1996 Feb;16(2):294-303. doi:10.1161/ 01.atv.16.2.294. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/8620346)
- Reshetnyak Y, Tchedre KT, Nair MP, Pritchard PH, Lacko AG. Structuraldifferences between wild-type and fish eye disease mutant of lecithin:cholesterolacyltransferase. J Biomol Struct Dyn. 2006 Aug;24(1):75-82. doi:10.1080/07391102.2006.10507101. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/16780378)
- Savel J, Lafitte M, Pucheu Y, Pradeau V, Tabarin A, Couffinhal T. Very lowlevels of HDL cholesterol and atherosclerosis, a variable relationship--a reviewof LCAT deficiency. Vasc Health Risk Manag. 2012;8:357-61. doi:10.2147/VHRM.S29985. Epub 2012 Jun 5. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/22701329) or Free article on PubMed Central (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3 373316/)
- Winder AF, Owen JS, Pritchard PH, Lloyd-Jones D, Vallance DT, White P, Wray R. A first British case of fish-eye disease presenting at age 75 years: a doubleheterozygote for defined and new mutations affecting LCAT structure and expression. J Clin Pathol. 1999 Mar;52(3):228-30. doi: 10.1136/jcp.52.3.228. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/10450185) or Free article on PubMed Central (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC501085/)

Last updated August 1, 2013