

Mitochondrial membrane protein-associated neurodegeneration

Description

Mitochondrial membrane protein-associated neurodegeneration (MPAN) is a disorder of the nervous system. The condition typically begins in childhood or early adulthood and worsens (progresses) over time.

MPAN commonly begins with difficulty walking. As the condition progresses, affected individuals usually develop other movement problems, including muscle stiffness (spasticity) and involuntary muscle cramping (dystonia). Many people with MPAN have a pattern of movement abnormalities known as parkinsonism. These abnormalities include unusually slow movement (bradykinesia), muscle rigidity, involuntary trembling (tremors), and an inability to hold the body upright and balanced (postural instability).

Other neurological problems that occur in individuals with MPAN include degeneration of the nerve cells that carry visual information from the eyes to the brain (optic atrophy), which can impair vision; problems with speech (dysarthria); difficulty swallowing (dysphagia); and, in later stages of the condition, an inability to control the bowels or the flow of urine (incontinence). Additionally, affected individuals may experience a loss of intellectual function (dementia) and psychiatric symptoms such as behavioral problems, mood swings, hyperactivity, and depression.

MPAN is characterized by an abnormal buildup of iron in certain regions of the brain. Because of these deposits, MPAN is considered part of a group of conditions known as neurodegeneration with brain iron accumulation (NBIA).

Frequency

MPAN is a rare condition that is estimated to affect less than 1 in 1 million people.

Causes

Mutations in the *C19orf12* gene cause MPAN. The protein produced from this gene is found in the membrane of cellular structures called mitochondria, which are the energy-producing centers of the cell. Although its function is unknown, researchers suggest that the C19orf12 protein plays a role in the maintenance of fat (lipid) molecules, a process known as lipid homeostasis.

The gene mutations that cause this condition lead to an altered C19orf12 protein that

likely has little or no function. It is unclear how these genetic changes lead to the neurological problems associated with MPAN. Researchers are working to determine whether there is a link between problems with lipid homeostasis and brain iron accumulation and how these abnormalities might contribute to the features of this disorder.

[Learn more about the gene associated with Mitochondrial membrane protein-associated neurodegeneration](#)

- C19orf12

Inheritance

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- Mitochondrial membrane protein-associated neurodegeneration due to C19orf12 mutation
- Mitochondrial protein-associated neurodegeneration
- MPAN
- NBIA4
- Neurodegeneration with brain iron accumulation 4

Additional Information & Resources

Genetic Testing Information

- Genetic Testing Registry: Neurodegeneration with brain iron accumulation 4 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C3280371/>)

Genetic and Rare Diseases Information Center

- Mitochondrial membrane protein-associated neurodegeneration (<https://rarediseases.info.nih.gov/diseases/12569/index>)

Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD) (<https://rarediseases.org/>)

Clinical Trials

- ClinicalTrials.gov (<https://clinicaltrials.gov/search?cond=%22Mitochondrial membrane protein-associated neurodegeneration%22>)

Catalog of Genes and Diseases from OMIM

- NEURODEGENERATION WITH BRAIN IRON ACCUMULATION 4; NBIA4 (<https://omim.org/entry/614298>)

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28mitochondrial+membrane+protein-associated+neurodegeneration%5BTIAB%5D%29+OR+%28nbia4%5BTIAB%5D%29+OR+%28mitochondrial+membrane+protein+associated+neurodegeneration%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D%29%29%29>)

References

- Dogu O, Krebs CE, Kaleagasi H, Demirtas Z, Oksuz N, Walker RH, Paisan-Ruiz C. Rapid disease progression in adult-onset mitochondrial membraneprotein-associated neurodegeneration. *Clin Genet*. 2013 Oct;84(4):350-5. doi:10.1111/cge.12079. Epub 2013 Jan 21. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23278385>)
- Gregory A, Klopstock T, Kmiec T, Hogarth P, Hayflick SJ. MitochondrialMembrane Protein-Associated Neurodegeneration. 2014 Feb 27 [updated 2021 Mar 4]. In: Adam MP, Bick S, Mirzaa GM, Pagon RA, Wallace SE, Amemiya A, editors. *GeneReviews* (R) [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2025. Available from <http://www.ncbi.nlm.nih.gov/books/NBK185329/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/24575447>)
- Hartig M, Prokisch H, Meitinger T, Klopstock T. Mitochondrial membraneprotein-associated neurodegeneration (MPAN). *Int Rev Neurobiol*. 2013;110:73-84.doi: 10.1016/B978-0-12-410502-7.00004-1. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/24209434>)
- Hartig MB, Iuso A, Haack T, Kmiec T, Jurkiewicz E, Heim K, Roeber S, Tarabin V, Dusi S, Krajewska-Walasek M, Jozwiak S, Hempel M, Winkelmann J, Elstner M, Oexle K, Klopstock T, Mueller-Felber W, Gasser T, Trenkwalder C, Tiranti V, Kretzschmar H, Schmitz G, Strom TM, Meitinger T, Prokisch H. Absence of an orphanmitochondrial protein, c19orf12, causes a distinct clinical subtype ofneurodegeneration with brain iron accumulation. *Am J Hum Genet*. 2011 Oct7;89(4):543-50. doi: 10.1016/j.ajhg.2011.09.007. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21981780>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3188837/>)

- Hogarth P, Gregory A, Kruer MC, Sanford L, Wagoner W, Natowicz MR, Egel RT, Subramony SH, Goldman JG, Berry-Kravis E, Foulds NC, Hammans SR, Desguerre I, Rodriguez D, Wilson C, Diedrich A, Green S, Tran H, Reese L, Woltjer RL, Hayflick SJ. New NBIA subtype: genetic, clinical, pathologic, and radiographic features of MPAN. *Neurology*. 2013 Jan 15;80(3):268-75. doi: 10.1212/WNL.0b013e31827e07be. Epub 2012 Dec 26. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23269600>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3589182/>)
- Schulte EC, Claussen MC, Jochim A, Haack T, Hartig M, Hempel M, Prokisch H, Haun-Junger U, Winkelmann J, Hemmer B, Forschler A, Ilg R. Mitochondrial membraneprotein associated neurodegeneration: a novel variant of neurodegeneration with brain iron accumulation. *Mov Disord*. 2013 Feb;28(2):224-7. doi:10.1002/mds.25256. Epub 2012 Nov 19. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23436634>)

Last updated January 1, 2017