

# CYP1B1 gene

cytochrome P450 family 1 subfamily B member 1

#### **Normal Function**

The CYP1B1 gene provides instructions for producing an enzyme that is a member of the cytochrome P450 family of enzymes. These enzymes are involved in many processes in the body, such as assisting with reactions that break down drugs and produce certain fats (lipids). The CYP1B1 enzyme participates in biochemical reactions in which an oxygen atom is added to other molecules.

The CYP1B1 enzyme is active in many tissues, including structures of the eye. The function of the CYP1B1 enzyme in the development of the eye is unclear, but it may play a role in forming structures at the front of the eye and may also be involved in a process that regulates the secretion of fluid inside the eye.

# **Health Conditions Related to Genetic Changes**

## Early-onset glaucoma

More than 140 *CYP1B1* gene variants (also called mutations) have been identified to cause early-onset glaucoma. People with this condition experience increased pressure within the eye before the age of 40. This pressure causes damage to the optic nerves connecting the eyes and the brain. Between 20 percent and 40 percent of people with glaucoma before the age of 3 (primary congenital glaucoma) have variants in the *CYP1B1* gene. The variants that cause early-onset glaucoma may result in an enzyme that is unstable or the wrong shape.

It is not well understood how changes in the CYP1B1 enzyme cause signs and symptoms of glaucoma. Recent studies suggest that the defects may interfere with the early development of the trabecular meshwork, which is a network of mesh-like canals that helps drain excess fluid from the eye. If fluid cannot be drained, pressure inside the eye may increase, which is characteristic of glaucoma.

#### Peters anomaly

A few variants in the *CYP1B1* gene have been found to cause Peters anomaly. This condition is characterized by abnormal development of certain structures at the front of the eye and clouding of the clear front surface of the eye (cornea). The variants that

cause Peters anomaly likely impair the normal function of the CYP1B1 enzyme or disrupt enzyme production. As a result, there is a shortage (deficiency) of normal enzyme. A lack of the CYP1B1 enzyme likely disrupts normal development of the eye, although it is unclear exactly how this deficiency leads to the features of Peters anomaly.

### Other Names for This Gene

- aryl hydrocarbon hydroxylase
- CP1B
- CP1B1\_HUMAN
- cytochrome P450, family 1, subfamily B, polypeptide 1
- cytochrome P450, subfamily I (dioxin-inducible), polypeptide 1 (glaucoma 3, primary infantile)
- flavoprotein-linked monooxygenase
- GLC3A
- microsomal monooxygenase
- xenobiotic monooxygenase

### **Additional Information & Resources**

## Tests Listed in the Genetic Testing Registry

Tests of CYP1B1 (https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=1545[geneid])

## Scientific Articles on PubMed

PubMed (https://pubmed.ncbi.nlm.nih.gov/?term=(CYP1B1%5BTIAB%5D)+AND+((Genes%5BMH%5D)+OR+(Genetic+Phenomena%5BMH%5D))+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D)

## Catalog of Genes and Diseases from OMIM

CYTOCHROME P450, SUBFAMILY I, POLYPEPTIDE 1; CYP1B1 (https://omim.org/entry/601771)

## Gene and Variant Databases

- NCBI Gene (https://www.ncbi.nlm.nih.gov/gene/1545)
- ClinVar (https://www.ncbi.nlm.nih.gov/clinvar?term=CYP1B1[gene])

### References

- Achary MS, Reddy AB, Chakrabarti S, Panicker SG, Mandal AK, Ahmed N, Balasubramanian D, Hasnain SE, Nagarajaram HA. Disease-causing mutations inproteins: structural analysis of the CYP1B1 mutations causing primary congenitalglaucoma in humans. Biophys J. 2006 Dec 15;91(12):4329-39. doi:10. 1529/biophysj.106.085498. Epub 2006 Sep 8. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/16963504) or Free article on PubMed Central (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1779944/)
- Chavarria-Soley G, Sticht H, Aklillu E, Ingelman-Sundberg M, Pasutto F, ReisA, Rautenstrauss B. Mutations in CYP1B1 cause primary congenital glaucoma byreduction of either activity or abundance of the enzyme. Hum Mutat. 2008Sep;29( 9):1147-53. doi: 10.1002/humu.20786. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/18470941)
- Chen Y, Jiang D, Yu L, Katz B, Zhang K, Wan B, Sun X. CYP1B1 and MYOCmutations in 116 Chinese patients with primary congenital glaucoma. ArchOphthalmol. 2008 Oct;126(10):1443-7. doi: 10.1001/archopht.126.10.1443. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/18852424)
- Melki R, Colomb E, Lefort N, Brezin AP, Garchon HJ. CYP1B1 mutations in Frenchpatients with early-onset primary open-angle glaucoma. J Med Genet. 2004Sep;41(9):647-51. doi: 10.1136/jmg.2004.020024. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/15342693) or Free article on PubMed Central (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1735887/)
- Selvan H, Gupta S, Wiggs JL, Gupta V. Juvenile-onset open-angle glaucoma -Aclinical and genetic update. Surv Ophthalmol. 2022 Jul-Aug;67(4):1099-1117. doi: 10.1016/j.survophthal.2021.09.001. Epub 2021 Sep 16. Citation on PubMed (https://www.ncbi.nlm.nih.gov/pubmed/34536459)
- Vasiliou V, Gonzalez FJ. Role of CYP1B1 in glaucoma. Annu Rev PharmacolToxicol. 2008;48:333-58. doi: 10.1146/annurev.pharmtox.48.061807. 154729. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/17914928)
- Vincent A, Billingsley G, Priston M, Glaser T, Oliver E, Walter M, Ritch R, Levin A, Heon E. Further support of the role of CYP1B1 in patients with Petersanomaly. Mol Vis. 2006 May 16;12:506-10. Citation on PubMed (https://pubmed.ncbi.nlm.nih.gov/1 6735991)

## **Genomic Location**

The CYP1B1 gene is found on chromosome 2 (https://medlineplus.gov/genetics/chromosome/2/).

Last updated April 4, 2022