

HLA-DQA1 gene

major histocompatibility complex, class II, DQ alpha 1

Normal Function

The *HLA-DQA1* gene provides instructions for making a protein that plays a critical role in the immune system. The *HLA-DQA1* gene is part of a family of genes called the human leukocyte antigen (HLA) complex. The HLA complex helps the immune system distinguish the body's own proteins from proteins made by foreign invaders such as viruses and bacteria.

The HLA complex is the human version of the major histocompatibility complex (MHC), a gene family that occurs in many species. The *HLA-DQA1* gene belongs to a group of MHC genes called MHC class II. MHC class II genes provide instructions for making proteins that are present on the surface of certain immune system cells. These proteins attach to protein fragments (peptides) outside the cell. MHC class II proteins display these peptides to the immune system. If the immune system recognizes the peptides as foreign (such as viral or bacterial peptides), it triggers a response to attack the invading viruses or bacteria.

The protein produced from the *HLA-DQA1* gene attaches (binds) to the protein produced from another MHC class II gene, *HLA-DQB1*. Together, they form a functional protein complex called an antigen-binding DQ $\alpha\beta$ heterodimer. This complex displays foreign peptides to the immune system to trigger the body's immune response.

Each MHC class II gene has many possible variations, allowing the immune system to react to a wide range of foreign invaders. Researchers have identified hundreds of different versions (alleles) of the *HLA-DQA1* gene, each of which is given a particular number (such as *HLA-DQA1*05:01*).

Health Conditions Related to Genetic Changes

Celiac disease

At least two specific combinations of HLA gene variants (HLA haplotypes) have been found to increase the risk of developing celiac disease, a disorder in which inflammation damages the intestinal tract and other organs and tissues. One of these haplotypes, known as DQ2, is composed of the protein produced from *HLA-DQA1* gene variants known as *HLA-DQA1*05:01* or *HLA-DQA1*05:05* bound to the protein produced from

HLA-DQB1 gene variants known as *HLA-DQB1*02:01* or *HLA-DQB1*02:02*. The other haplotype, known as DQ8, is composed of the protein produced from *HLA-DQA1* gene variants known as *HLA-DQA1*03:01* or *HLA-DQA1*03:02* bound to the protein produced from the *HLA-DQB1* gene variant known as *HLA-DQB1*03:02*.

The DQ2 and DQ8 haplotypes, which may occur separately or together, seem to increase the risk of an inappropriate immune response to the protein gluten, which is found in wheat, rye, and barley. This immune system malfunction results in the damage to the body's organs and tissues that occurs in celiac disease. However, the DQ2 and DQ8 haplotypes are also found in 30 percent of the general population, and only 3 percent of individuals with these haplotypes develop celiac disease.

Alopecia areata

MedlinePlus Genetics provides information about Alopecia areata

Autoimmune Addison disease

MedlinePlus Genetics provides information about Autoimmune Addison disease

Idiopathic inflammatory myopathy

MedlinePlus Genetics provides information about Idiopathic inflammatory myopathy

Juvenile idiopathic arthritis

MedlinePlus Genetics provides information about Juvenile idiopathic arthritis

Narcolepsy

MedlinePlus Genetics provides information about Narcolepsy

Rosacea

MedlinePlus Genetics provides information about Rosacea

Type 1 diabetes

Combinations of variations in the *HLA-DQA1* gene and other HLA genes affect the risk of type 1 diabetes. Type 1 diabetes is characterized by high blood sugar (glucose) levels resulting from a shortage of the hormone insulin and is caused by autoimmune damage to insulin-producing cells in the pancreas.

Type 1 diabetes risk is most increased by two HLA haplotypes involving variations of the *HLA-DQA1* and *HLA-DQB1* genes and another HLA gene called *HLA-DRB1*. One haplotype, written as *DRB1*03:01-DQA1*05:01-DQB1*02*, is called DR3. The other haplotype, written as *DRB1*04:01/02/04/05/08-DQA1*03:01-DQB1*02*, is called DR4. People at highest risk of developing type 1 diabetes have one copy of the DR3 haplotype and one copy of the DR4 haplotype in each cell. Other HLA haplotypes only

mildly increase the risk of type 1 diabetes, while some haplotypes seem to protect against developing this condition. Variations in other genes and environmental factors are also thought to affect the risk of this complex disorder.

Autoimmune disorders

Certain normal variations of the *HLA-DQA1* gene have been associated with increased risk of autoimmune disorders, which occur when the immune system malfunctions and attacks the body's own tissues and organs. It is unclear how different versions of the *HLA-DQA1* gene influence the risk of developing autoimmune disorders. These conditions are thought to result from a combination of multiple environmental and genetic factors. Changes in other HLA and non-HLA genes, some of which remain unknown, also likely contribute to the risk of developing these complex conditions.

Other disorders

Normal variations in the *HLA-DQA1* gene can affect the body's ability to recognize and react to foreign invaders (pathogens). For example, variations of this gene have been shown to increase or decrease a person's chance of getting infections such as hepatitis B and leprosy or may affect the severity of illness if infection occurs.

A particular variant of the *HLA-DQA1* gene known as *HLA-DQA1*02:01* increases the risk of liver damage in women with advanced breast cancer treated with a drug called lapatinib. Researchers suggest that the variant may increase immune system sensitivity to the drug, resulting in inflammation that damages the liver.

Other Names for This Gene

- DC-1 alpha chain
- DC-alpha
- DQ-A1
- FLJ27088
- FLJ27328
- GSE
- HLA class II histocompatibility antigen, DQ alpha 1 chain
- HLA class II histocompatibility antigen, DQ alpha 1 chain precursor
- HLA class II histocompatibility antigen, DQ(W3) alpha chain
- HLA-DCA
- HLA-DQA
- leucocyte antigen DQA1
- leukocyte antigen alpha chain
- MGC149527
- MHC class II antigen
- MHC class II DQA1

- MHC class II HLA-D alpha glycoprotein
- MHC class II HLA-DQ-alpha-1
- MHC class II surface glycoprotein
- MHC HLA-DQ alpha

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of HLA-DQA1 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=3117\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=3117[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28HLA-DQA1%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>)

Catalog of Genes and Diseases from OMIM

- MAJOR HISTOCOMPATIBILITY COMPLEX, CLASS II, DQ ALPHA-1; HLA-DQA1 (<https://omim.org/entry/146880>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/3117>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=HLA-DQA1\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=HLA-DQA1[gene]))

References

- Abadie V, Sollid LM, Barreiro LB, Jabri B. Integration of genetic and immunological insights into a model of celiac disease pathogenesis. *Annu Rev Immunol.* 2011;29:493-525. doi: 10.1146/annurev-immunol-040210-092915. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21219178>)
- da Silva SA, Mazini PS, Reis PG, Sell AM, Tsuneto LT, Peixoto PR, Visentainer JE. HLA-DR and HLA-DQ alleles in patients from the south of Brazil: markers for leprosy susceptibility and resistance. *BMC Infect Dis.* 2009 Aug 22;9:134. doi:10.1186/1471-2334-9-134. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19698125>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2746224/>)
- Lipton RB, Drum M, Greeley SA, Danielson KK, Bell GI, Hagopian WA. HLA-DQ haplotypes differ by ethnicity in patients with childhood-onset diabetes.

PediatrDiabetes. 2011 Jun;12(4 Pt 2):388-95. doi: 10.1111/j.1399-5448.2010.00712.x. Epub 2011 Mar 21. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21418452>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3406606/>)

- Noble JA, Valdes AM. Genetics of the HLA region in the prediction of type 1 diabetes. *Curr Diab Rep*. 2011 Dec;11(6):533-42. doi: 10.1007/s11892-011-0223-x. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21912932>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3233362/>)
- Spraggs CF, Budde LR, Briley LP, Bing N, Cox CJ, King KS, Whittaker JC, Mooser VE, Preston AJ, Stein SH, Cardon LR. HLA-DQA1*02:01 is a major risk factor for lapatinib-induced hepatotoxicity in women with advanced breast cancer. *J Clin Oncol*. 2011 Feb 20;29(6):667-73. doi: 10.1200/JCO.2010.31.3197. Epub 2011 Jan 18. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21245432>)
- Taylor AK, Lebowitz B, Snyder CL, Green PHR. Celiac Disease. 2008 Jul 3 [updated 2019 Jan 31]. In: Adam MP, Feldman J, Mirzazadeh GM, Pagon RA, Wallace SE, Amemiya A, editors. *GeneReviews*(R) [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2025. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1727/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301720>)
- Tjon JM, van Bergen J, Koning F. Celiac disease: how complicated can it get? *Immunogenetics*. 2010 Oct;62(10):641-51. doi: 10.1007/s00251-010-0465-9. Epub 2010 Jul 27. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20661732>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2944025/>)
- Uibo R, Tian Z, Gershwin ME. Celiac disease: a model disease for gene-environment interaction. *Cell Mol Immunol*. 2011 Mar;8(2):93-5. doi:10.1038/cmi.2010.62. Epub 2011 Feb 14. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21317918>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4003132/>)
- Xun YH, Guo JC, Shi WZ, Shi JP, Liu CL. [Association between HLA-DQA1 gene polymorphism and the outcomes of hepatitis B virus infection]. *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi*. 2009 Dec;23(6):430-3. Chinese. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20718347>)

Genomic Location

The *HLA-DQA1* gene is found on chromosome 6 (<https://medlineplus.gov/genetics/chromosome/6/>).

Last updated March 1, 2013