

## **STAT1 gene**

signal transducer and activator of transcription 1

### **Normal Function**

The *STAT1* gene provides instructions for making a protein that is involved in multiple immune system functions, including the body's defense against a fungus called *Candida*. When the immune system recognizes *Candida*, it generates cells called Th17 cells. These cells produce signaling molecules (cytokines) called the interleukin-17 (IL-17) family as part of an immune process called the IL-17 pathway. The IL-17 pathway creates inflammation, sending other cytokines and white blood cells that fight foreign invaders and promote tissue repair. In addition, the IL-17 pathway promotes the production of certain antimicrobial protein segments (peptides) that control growth of *Candida* on the surface of mucous membranes.

The STAT1 protein helps keep the immune system in balance by controlling the IL-17 pathway. When the STAT1 protein is turned on (activated), it blocks (inhibits) this pathway.

In contrast to its inhibitory role in the IL-17 pathway, the STAT1 protein helps promote other immune processes called the interferon-alpha/beta (IFNA/B) and interferon-gamma (IFNG) signaling pathways. The IFNA/B pathway is important in defense against viruses, and the IFNG pathway helps fight a type of bacteria called mycobacteria, which includes the bacterium that causes tuberculosis.

### **Health Conditions Related to Genetic Changes**

#### Familial candidiasis

At least 35 *STAT1* gene mutations have been identified in people with familial candidiasis, an inherited tendency to develop infections caused by the *Candida* fungus (commonly called yeast infections). Most people with familial candidiasis have chronic infections of the skin, nails, and mucous membranes such as the lining of the mouth, collectively called chronic mucocutaneous candidiasis, beginning in early childhood.

Some people with *STAT1* gene mutations have additional features such as increased susceptibility to other infections and an increased risk of autoimmune disorders, in which the immune system attacks the body's own tissues or organs. This combination of signs and symptoms caused by *STAT1* gene mutations is sometimes called

immunodeficiency 31C.

The *STAT1* gene mutations that have been identified in people with familial candidiasis are described as "gain of function" because they increase the amount of activated *STAT1* protein in cells. By increasing *STAT1*'s inhibitory effect on the IL-17 pathway, the mutations impair the body's ability to fight *Candida* and result in the chronic infection that occurs in familial candidiasis. The effects of the increase in activated *STAT1* protein on other signaling pathways are thought to underlie the variety of features that can occur in immunodeficiency 31C, but the specific mechanisms are not well understood.

### Shingles

MedlinePlus Genetics provides information about Shingles

### Other disorders

*STAT1* gene mutations can also cause other immune system problems called immunodeficiency 31A and immunodeficiency 31B. In contrast with the gain-of-function mutations associated with familial candidiasis (described above), these *STAT1* gene mutations are described as "loss-of-function" mutations because they impair or eliminate the protein's normal action.

The *STAT1* gene mutations that cause immunodeficiency 31A impair the *STAT1* protein's function in the IFNG pathway, resulting in an increased susceptibility to mycobacteria. In the more severe disorder immunodeficiency 31B, the *STAT1* gene mutations impair both the IFNG pathway and the IFNA/B pathway, resulting in susceptibility to both viral infections such as herpes simplex and viral encephalitis, and mycobacterial infections such as tuberculosis.

### **Other Names for This Gene**

- IMD31A
- IMD31B
- IMD31C
- ISGF-3
- signal transducer and activator of transcription 1, 91kDa
- signal transducer and activator of transcription-1
- STAT91
- transcription factor ISGF-3 components p91/p84

### **Additional Information & Resources**

#### Tests Listed in the Genetic Testing Registry

- Tests of *STAT1* ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=6772\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=6772[geneid]))

## Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28STAT1%5BTI%5D%29+OR+%28signal+transducer+and+activator+of+transcription+1%5BTI%5D%29%29+AND+%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D>)

## Catalog of Genes and Diseases from OMIM

- SIGNAL TRANSDUCER AND ACTIVATOR OF TRANSCRIPTION 1; STAT1 (<https://omim.org/entry/600555>)
- IMMUNODEFICIENCY 31A; IMD31A (<https://omim.org/entry/614892>)
- IMMUNODEFICIENCY 31B; IMD31B (<https://omim.org/entry/613796>)

## Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/6772>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=STAT1\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=STAT1[gene]))

## **References**

- Boisson-Dupuis S, Kong XF, Okada S, Cypowyj S, Puel A, Abel L, Casanova JL. Inborn errors of human STAT1: allelic heterogeneity governs the diversity of immunological and infectious phenotypes. *Curr Opin Immunol.* 2012 Aug;24(4):364-78. doi: 10.1016/j.co.2012.04.011. Epub 2012 May 30. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22651901>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3477860/>)
- Depner M, Fuchs S, Raabe J, Frede N, Glocker C, Doffinger R, Gkrania-Klotsas E, Kumararatne D, Atkinson TP, Schroeder HW Jr, Niehues T, Duckers G, Stray-Pedersen A, Baumann U, Schmidt R, Franco JL, Orrego J, Ben-Shoshan M, McCusker C, Jacob CM, Carneiro-Sampaio M, Devlin LA, Edgar JD, Henderson P, Russell RK, Skytte AB, Seneviratne SL, Wanders J, Stauss H, Meyts I, Moens L, Jesenak M, Kobbe R, Borte S, Borte M, Wright DA, Hagin D, Torgerson TR, Grimbacher B. The Extended Clinical Phenotype of 26 Patients with Chronic Mucocutaneous Candidiasis due to Gain-of-Function Mutations in STAT1. *J Clin Immunol.* 2016 Jan;36(1):73-84. doi: 10.1007/s10875-015-0214-9. Epub 2015 Nov 25. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/26604104>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4718942/>)
- Liu L, Okada S, Kong XF, Kreins AY, Cypowyj S, Abhyankar A, Toubiana J, Itan Y, Audry M, Nitschke P, Masson C, Toth B, Flatot J, Migaud M, Chrabieh M, Kochetkov

- T, Bolze A, Borghesi A, Toulon A, Hiller J, Eyerich S, Eyerich K, Gulacsy V, Chernyshova L, Chernyshov V, Bondarenko A, Grimaldo RM, Blancas-Galicia L, Beas IM, Roesler J, Magdorf K, Engelhard D, Thumerelle C, Burgel PR, Hoernes M, Drexel B, Seger R, Kusuma T, Jansson AF, Sawalle-Belohradsky J, Belohradsky B, Jouanguy E, Bustamante J, Bue M, Karin N, Wildbaum G, Bodemer C, Lortholary O, Fischer A, Blanche S, Al-Muhsen S, Reichenbach J, Kobayashi M, Rosales FE, Lozano CT, Kilic SS, Oleastro M, Etzioni A, Traidl-Hoffmann C, Renner ED, Abel L, Picard C, Marodi L, Boisson-Dupuis S, Puel A, Casanova JL. Gain-of-function human STAT1 mutations impair IL-17 immunity and underlie chronic mucocutaneous candidiasis. *J Exp Med.* 2011 Aug 1;208(8):1635-48. doi: 10.1084/jem.20110958. Epub 2011 Jul 4. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21727188/>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3149226/>)
- Mizoguchi Y, Tsumura M, Okada S, Hirata O, Minegishi S, Imai K, Hyakuna N, Muramatsu H, Kojima S, Ozaki Y, Imai T, Takeda S, Okazaki T, Ito T, Yasunaga S, Takihara Y, Bryant VL, Kong XF, Cypowij S, Boisson-Dupuis S, Puel A, Casanova JL, Morio T, Kobayashi M. Simple diagnosis of STAT1 gain-of-function alleles in patients with chronic mucocutaneous candidiasis. *J Leukoc Biol.* 2014 Apr;95(4):667-76. doi: 10.1189/jlb.0513250. Epub 2013 Dec 16. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/24343863/>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3958742/>)
  - Smeekens SP, van de Veerdonk FL, Kullberg BJ, Netea MG. Genetic susceptibility to *Candida* infections. *EMBO Mol Med.* 2013 Jun;5(6):805-13. doi: 10.1002/emmm.201201678. Epub 2013 Apr 30. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23629947/>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3779444/>)
  - Soltesz B, Toth B, Shabashova N, Bondarenko A, Okada S, Cypowij S, Abhyankar A, Csorba G, Tasko S, Sarkadi AK, Mehes L, Rozsival P, Neumann D, Chernyshova L, Tulassay Z, Puel A, Casanova JL, Sediva A, Litzman J, Marodi L. New and recurrent gain-of-function STAT1 mutations in patients with chronic mucocutaneous candidiasis from Eastern and Central Europe. *J Med Genet.* 2013 Sep;50(9):567-78. doi: 10.1136/jmedgenet-2013-101570. Epub 2013 May 24. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23709754/>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3756505/>)
  - Toubiana J, Okada S, Hiller J, Oleastro M, Lagos Gomez M, Aldave Becerra JC, Ouachee-Chardin M, Fouyssac F, Girisha KM, Etzioni A, Van Montfrans J, Camcioglu Y, Kerns LA, Belohradsky B, Blanche S, Bousfiha A, Rodriguez-Gallego C, Meyts I, Kisand K, Reichenbach J, Renner ED, Rosenzweig S, Grimbacher B, van de Veerdonk FL, Traidl-Hoffmann C, Picard C, Marodi L, Morio T, Kobayashi M, Lilic D, Milner JD, Holland S, Casanova JL, Puel A; International STAT1 Gain-of-Function Study Group. Heterozygous STAT1 gain-of-function mutations underlie an unexpectedly broad clinical phenotype. *Blood.* 2016 Jun 23;127(25):3154-64. doi: 10.1182/blood-2015-11-679902. Epub 2016 Apr 25. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/27114460/>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4920021/>)
  - Yamazaki Y, Yamada M, Kawai T, Morio T, Onodera M, Ueki M, Watanabe N,

Takada H, Takezaki S, Chida N, Kobayashi I, Ariga T. Two novel gain-of-function mutations of STAT1 responsible for chronic mucocutaneous candidiasis disease: impaired production of IL-17A and IL-22, and the presence of anti-IL-17F autoantibody. *J Immunol*. 2014 Nov 15;193(10):4880-7. doi:10.4049/jimmunol.1401467. Epub 2014 Oct 6. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/25288569>)

## Genomic Location

The *STAT1* gene is found on chromosome 2 (<https://medlineplus.gov/genetics/chromosome/2/>).

**Last updated September 1, 2016**